RESTRICTED USE PESTICIDE Toxic to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only those uses covered by the certified applicator's certification.



FOR CONTROL OF INSECT PESTS ON CANEBERRIES

EPA Registration No. 279-3069

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CROP	PEST	RATE OF APPLICATION
Caneberries	Raspberry Crown Borer	6.4 oz product/acre (0.1 lbs a.i./acre)

Directions for use:

Apply 0.1 lb ai / a, post-harvest (fall) or pre-bloom (spring), as a drench application directed at the crown of plants in a minimum of 200 gallons water / acre. Greater efficacy is observed at higher water gallonages (up to 400 gallons/a) or in an application prior to a significant rainfall event. Do not make pre-bloom foliar and pre-bloom drench applications.

RESTRICTIONS:

Do not apply more than 0.20 lbs. a.i. per acre per season

Do not apply within 3 days of harvest.

KEEP OUT OF REACH OF CHILDREN WARNING

For Statement of Practical Treatment and other Precautionary Statements, see label attached to container.

All applicable directions, restrictions, Worker Protection Standard requirements and other precautions on the EPA-registered label are to be followed.

NOTE: This label must be in the possession of the user at the time of pesticide application.

FMC Corporation, Agricultural Products Group, Philadelphia, PA 19103