

Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

Provado® 1.6 Flowable Insecticide

**EPA Reg. No. 264-763
(formerly EPA Reg. No. 3125-457)**

Approval: June 18, 2003

Label Revisions: *Added use directions for Stone Fruit. Revised protective glove requirement. Revised Rotational Crop instructions.*

Supplemental Label

THIS SUPPLEMENTAL LABEL IS NOT FOR USE IN CALIFORNIA, FLORIDA, HAWAII OR NEW YORK.

Provado® 1.6 Flowable Insecticide

For control of certain insects infesting various crops.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read this label and the product package label before using this product. This Supplemental Label must be in possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the registered product label for Provado® 1.6 Flowable Insecticide.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing.

Applicators and Other Handlers Must Wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.
- Shoes plus socks

Follow manufacturer's instructions for cleaning/ maintaining personal protective equipment, PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application.

For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval should be observed.

ROTATIONAL PLANT-BACK INTERVALS¹			
IMMEDIATE PLANT-BACK (all crops on this label plus the following crops not on this label)			
Barley	Corn (field, sweet & pop)	Florence fennel	Rhubarb
Canola	Celtuce	Leafy petioles	Sorghum
Cardoon	Cranberry	Mustard Seed	Sugarbeets
Chinese celery	Cucurbits	Rape seed	Swiss chard
			Wheat
30-DAY PLANT-BACK			
Cereals (including buckwheat, millet, oats, rice, rye, and triticale)	Soybeans	Safflower	
12-MONTH PLANT-BACK			
All Other Crops			

¹ Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed.

RECOMMENDED APPLICATIONS

STONE FRUIT			
CROP	PEST	Rate fluid ounces/100 gallons	Rate fluid ounces/Acre
Apricot Nectarine Peach (For other stone fruit crops, see below)	Aphids	2.0	4.0 – 8.0
	Green June beetle		
	Japanese beetle		
	Potato leafhopper		
	Rose chafer		
	San Jose scale		
	Tarnished plant bug		
	White apple leafhopper		
	Pests Suppressed	2.0	8.0
	Plum curculio		
	Stink bugs		

NOTES AND RESTRICTIONS:

Pre-Harvest Interval (PHI): **0 days**

Maximum Provado® allowed per 7-day interval: **8.0 fluid ounces/Acre** (0.10 lbs AI/A).

Maximum Provado® allowed per crop season: **24.0 fluid ounces/Acre** (0.30 lbs AI/A).

Minimum application volume (water): 50 GPA – ground application, 25 GPA – aerial application.

Maximum concentrate rate is based on canopy size requiring 400 gallons per acre, if sprayed to drip.

For concentrate spray application based on tree row-volume/density, do not apply less than 4.0 fluid ounces/A.

APPLICATIONS:

Apply specified rate per acre as foliar spray with ground or aerial equipment after pollination is complete and bees are no longer present in the orchard. Thorough, uniform coverage of fruit and foliage is necessary to achieve optimal control.

Applications targeting San Jose Scale should be timed to the crawler stage. Treat each generation.

The amount of Provado® required per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees. For example, to calculate the rate for aphid control on smaller trees, multiply 2 fluid ounces times the number of 100 gallons of spray solution required to thoroughly wet, just prior to the point of runoff, one acre of the trees being treated. For concentrate sprays, apply the same amount of product per acre as would be applied in a dilute spray based on tree size and foliage volume.

CROP	PEST	Rate fluid ounces/100 gallons	Rate fluid ounces/Acre
Cherry (sweet and tart) Plum (including Chickasaw, Damson and Japanese) Plumcot Prune (For other stone fruit crops, see above)	Aphids	2.0	4.0 – 8.0
	Green June beetle		
	Japanese beetle		
	Potato leafhopper		
	Rose chafer		
	San Jose scale		
	Tarnished plant bug		
	Western cherry fruit fly		
	White apple leafhopper		
	Pests Suppressed	2.0	8.0
	Plum curculio		
	Stink bugs		

NOTES AND RESTRICTIONS:

Pre-Harvest Interval (PHI): **7 days**

Maximum Provado® allowed per 10-day interval: **8.0 fluid ounces/Acre** (0.10 lbs AI/A).

Maximum Provado® allowed per crop season: **40.0 fluid ounces/Acre** (0.50 lbs AI/A).

Minimum application volume (water): 50 GPA – ground application, 25 GPA – aerial application.

Maximum concentrate rate is based on canopy size requiring 400 gallons per acre, if sprayed to drip.

For concentrate spray application based on tree row-volume/density, do not apply less than 4.0 fluid ounces/A.

APPLICATIONS:

Apply specified rate per acre as foliar spray with ground or aerial equipment after pollination is complete and bees are no longer present in the orchard. Thorough, uniform coverage of fruit and foliage is necessary to achieve optimal control.

Applications targeting San Jose Scale should be timed to the crawler stage. Treat each generation.

Applications targeting fruit fly should begin at adult emergence and continue on a 10 day spray interval through the egg hatch period. Proper application timing and full-labeled rates are critical for optimum performance on this pest.

The amount of Provado® required per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees. For example, to calculate the rate for aphid control on smaller trees, multiply 2 fluid ounces times the number of 100 gallons of spray solution required to thoroughly wet, just prior to the point of runoff, one acre of the trees being treated. For concentrate sprays, apply the same amount of product per acre as would be applied in a dilute spray based on tree size and foliage volume.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at www.bayercropscienceus.com

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

Provado is a registered trademark of Bayer.