

Supplemental Labeling
Post-bloom Application to Trunk for Borer Control
 (For Distribution and Use only in States East of the Rockies)

Lorsban^{*} 50-W

Insecticide

IN WATER SOLUBLE BAGS

EPA Reg. No. 62719-221-10163

ATTENTION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of application. Read the label affixed to the container for Lorsban^{*} 50W In Water Soluble Bags insecticide before applying. Carefully follow all precautionary statements and applicable use directions. Use of Lorsban 50W according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

Directions for Use
Use and Dosage Recommendations

Pests Controlled	Rate Lorsban 50W	Specific Use Directions
Dogwood Borer, American Plum Borer, Apple Bark Borer, Roundheaded Apple Tree Borer, Flatheaded Appletree Borer, Broad Necked Root Borer, Tilehoned Prionus	3 lb/100 gal of spray	Mix with water and apply directly to trunk from a distance of no more than 4 ft using low volume handgun or shielded spray equipment. Do not allow spray to contact foliage or fruit. Up to 2 applications may be made with a minimum spray interval of 14 days between applications.
		<ul style="list-style-type: none"> • Treat only the lower 4 feet of the trunk. • Do not make more than two applications per year for borer control. • Do not apply when wind speed is greater than 10 mph. • Do not apply within 28 days of harvest. • If the available residue data submitted in 2002 demonstrates the detection of chlorpyrifos at concentrations greater than 0.01 ppm, then Dow AgroSciences and EPA agree to discuss further mitigation.

*Trademark of DowAgroSciences, LLC
 ® Denotes Registered Trademark.